

**Zika and Spondweni viruses: Historic evidence of misidentification, misdiagnosis, and serious clinical disease manifestations**

Andrew D. Haddow<sup>1\*</sup> and John P. Woodall<sup>2</sup>

<sup>1</sup>United States Army Medical Research Institute of Infectious Diseases (USAMRIID)  
Entomology Department  
Fort Detrick, MD 21702-5011  
Andrew D. Haddow, Email: [andrew.d.haddow.ctr@mail.mil](mailto:andrew.d.haddow.ctr@mail.mil)

<sup>2</sup>ProMED-mail  
International Society for Infectious Diseases  
Brookline, MA, USA, 02446  
John P. Woodall, Email: [jackwoodall13@gmail.com](mailto:jackwoodall13@gmail.com)

\*Corresponding author

Content Type: Perspective

Word Count: 974

**Abstract**

The Spondweni serogroup (family *Flaviviridae*, genus *Flavivirus*) consists of two members: Zika and Spondweni viruses. Both viruses have been historically misidentified and their diseases have been misdiagnosed due to their serological cross-reactivity and similar clinical presentations. Within historic case reports a sub-set of patients has presented with short duration clinical manifestations suggestive of more serious illness.

## Perspective

Viruses within the genus *Flavivirus*, family *Flaviviridae*, are notorious for their serological cross-reactivity. Prior to the advent of genetic sequencing, classic serological assays such as the virus neutralization and hemagglutination-inhibition were used to differentiate the various species of arboviruses. Although not perfect, much of the early work differentiating flaviviruses into various serogroups was later confirmed by sequencing and phylogenetic analyses. Historically, serological assays (the neutralization and complement fixation tests) were also used to determine the prevalence of prior infection and geographic distribution of flaviviruses. Both viruses in the Spondweni serogroup, Zika (ZIKV) and Spondweni (SPONV), exhibit serological cross-reactivity and non-specific febrile illness, making diagnosis challenging in regions where both viruses co-circulate.

ZIKV was first isolated in Uganda in 1947 (strain MR-766) (1) and SPONV was first isolated in Nigeria in 1952 (strain Chuku) (2). Cross-reactivity using the neutralization test led to the misidentification of the SPONV Chuku strain isolated by MacNamara in 1952 as a strain of ZIKV (2-5). This misidentification led to additional studies where this strain of SPONV was reported as ZIKV – confusion that continues to the present day, although the misidentification of this isolate was clarified and widely reported in 1964 (3-5). Consequently, the clinical case reports by MacNamara (2), the work by Bearcroft (6) involving the experimental infection of a human volunteer and vector competence studies in *Aedes aegypti* mosquitoes, and the experimental work by Bearcroft (7) in *Macaca mulatta* monkeys investigating the effect of prior infection, and the subsequent histopathology of the liver and level of cross-protection following

47 exposure to yellow fever virus, all utilized SPONV (Chuku strain) rather than ZIKV.

48 Furthermore, early serosurveys in regions where both ZIKV and SPONV co-circulate are suspect  
49 due to serological cross-reactivity or where serological assays only screened for one of the two  
50 viruses (2, 8).

51  
52 Both ZIKV and SPONV are arthropod-borne viruses and utilize a mosquito/host (non-human  
53 primate and/or human) transmission cycle. ZIKV has a wide geographic distribution that  
54 includes East and West Africa, the Indian sub-continent, Southeast Asia, Oceania, South and  
55 Central America, and the Caribbean (9-11); whereas SPONV has thus far only been reported  
56 from sub-Saharan Africa (12). The lack of continuous historic detection in those regions with  
57 ZIKV or SPONV isolations or serological evidence of transmission prior to 2007 is likely due to  
58 the lack of surveillance, misdiagnosis, and under-reporting. It is plausible that virus outbreaks  
59 did occur in those regions with prior serologic evidence of infection and were attributed to other  
60 arboviral infections. Clinical ZIKV and SPONV presentation is similar to classic dengue fever  
61 which may have led to historic misdiagnosis (9).

62  
63 In their historic geographic ranges (distribution prior to 2007), both ZIKV and SPONV likely  
64 circulated at low levels in sylvatic cycles, whereby low numbers of naïve persons were  
65 periodically exposed to infection. While historic case reports of serious clinical manifestations  
66 associated with Spondweni Serogroup viruses may have been limited due to poor diagnosis and  
67 reporting, it is plausible that the lack of historic reports of congenital birth defects associated  
68 with ZIKV infection *in utero*, were a result of exposure to the virus prior to puberty. Such an  
69 infection would likely result in a female being immune to a subsequent infection during her

70 reproductive years. The extent of cross-protection exhibited within the Spondweni Serogroup is  
71 unknown, and cross-protection with other flaviviruses such as yellow fever virus and/or the 17D  
72 yellow fever virus vaccine appears to be limited (4-6, 13-16).

73  
74 Most cases of ZIKV and SPONV infection are asymptomatic (10, 11). Of symptomatic cases,  
75 signs and symptoms appear as early as 3 days following infection for both ZIKV(11) and  
76 SPONV (6). The common clinical presentation of ZIKV infection is now well established and a  
77 recent literature review lists the most common signs and symptoms reported from 195 patients  
78 from 1964 to 2016 as rash (67.2%), fever (63.6%), arthralgia (28.7%), myalgia (23.6%),  
79 headache (21.5%), conjunctivitis (20.5%), retro-orbital pain (11.3%), edema (9.7%), puritus  
80 (7.7%) and fatigue (7.2%) (11). Less is known regarding the clinical presentation of SPONV as  
81 there are few well documented clinical cases reported in the literature (n = 6). The most common  
82 signs and symptoms reported in those SPONV cases include: fever (100%), headache (83.3%),  
83 nausea (83.3%), myalgia (66.6%), arthralgia (50.0%), vertigo (33.3%), conjunctivitis (16.7%),  
84 macropapular and pruritic rash (16.7%), and epistaxis (16.7%), photophobia (16.7%), vomiting  
85 (16.7%), and disorientation (16.7%) (2, 5, 6, 13, 16). It is clear that both ZIKV and SPONV  
86 display similar signs and symptoms making diagnosis difficult in regions where both viruses co-  
87 circulate, additionally there are no commercially available serological assays that can  
88 differentiate these two viruses. In regions where both viruses co-circulate, diagnosis requires a  
89 monotypic reaction to a given serologic assay, PCR confirmation, or virus isolation.

90  
91 While most symptomatic ZIKV and SPONV infections present as a mild to moderate febrile  
92 illness, a sub-set of cases present with short duration clinical manifestations suggestive of more

serious illness. Prior to reports of ZIKV in the Western Hemisphere, Spondweni Serogroup case reports included: conjunctivitis (ZIKV and SPONV), macropapular rash (ZIKV and SPONV), pruritic rash (SPONV), hematuria (ZIKV), hematospermia (ZIKV), aphthous ulcer (ZIKV), and epistaxis (SPONV) indicating vascular leakage; and reports of photophobia (ZIKV and SPONV), vomiting (ZIKV and SPONV), vertigo (SPONV), disorientation (SPONV), meningismus (SPONV), and bilateral transient ocular paresis (SPONV) were indicative of neurological involvement (2, 5, 6, 13, 16, 17). Additionally, Guillain-Barré syndrome (18), evidence of sexual transmission (15), and evidence perinatal transmission (19) were all associated with a sub-set of ZIKV cases prior to the introduction of the virus in the Western Hemisphere.

In summary, symptomatic cases of ZIKV and SPONV infection present with similar signs and symptoms and anti-ZIKV and anti-SPONV serological assays exhibit cross-reactivity. Early work by MacNamara (1954) and Bearcroft (1956 and 1957) misidentified SPONV as ZIKV. Prior to the introduction of ZIKV into the Western Hemisphere there was evidence of clinical manifestations indicative of vascular leakage and neurological involvement within the Serogroup as well as unique transmission mechanisms associated with a sub-set of patients infected with ZIKV.

116 **Authors' contributions**

117

118 ADH and JPW both contributed equally to researching historic reports and writing the  
119 manuscript.

120

121 **Disclosure Statement**

122

123 The views expressed in this article are those of the authors and do not reflect the official policy  
124 or position of the U.S. Department of Defense or the Department of the Army.

125

126 **Competing interests**

127

128 The authors have no financial, personal, or professional interests that inappropriately influenced  
129 this paper.

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

**References**

1. Dick GW, Kitchen SF, Haddow AJ. Zika virus. I. Isolations and serological specificity. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1952;46(5):509-20.
2. Macnamara FN. Zika virus: a report on three cases of human infection during an epidemic of jaundice in Nigeria. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1954;48(2):139-45.
3. Haddow AJ, Williams MC, Woodall JP, Simpson DI, Goma LK. Twelve isolations of Zika virus from *Aedes* (*Stegomyia*) *africanus* (Theobald) taken in and above a Uganda Forest. Bulletin of the World Health Organization. 1964;31:57-69.
4. Simpson DI. Zika virus infection in man. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1964;58:335-8.
5. Draper CC. Infection with the Chuku strain of Spondweni virus. West African Medical Journal. 1965;14:16-9. Epub 1965/02/01.
6. Bearcroft WG. Zika virus infection experimentally induced in a human volunteer. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1956;50(5):442-8.
7. Bearcroft WG. The histopathology of the liver of yellow fever-induced rhesus monkeys. The Journal of Pathology and Bacteriology. 1957;74:295-303.
8. Macnamara FN, Horn DW, Porterfield JS. Yellow fever and other arthropod-borne viruses: A consideration of two serological surveys made in South Western Nigeria. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1959;53(2):202-12.



- 167 9. Haddow AD, Schuh AJ, Yasuda CY, Kasper MR, Heang V, Huy R, et al. Genetic  
 168 characterization of Zika virus strains: Geographic expansion of the Asian lineage. PLoS  
 169 Neglected Tropical Diseases. 2012;6(2):e1477.
- 170 10. Petersen LR, Jamieson DJ, Powers AM, Honein MA. Zika Virus. The New England  
 171 Journal of Medicine. 2016; 374:1552-1563
- 172 11. Plourde AR, Bloch EV. A Literature Review of Zika Virus. Emerging Infectious  
 173 Diseases. 2016;22(7).
- 174 12. Theiler M, Downs WG. The arthropod-borne viruses of vertebrates. New Haven and  
 175 London: Yale University Press; 1973.
- 176 13. McIntosh BM, Kokernot RH, Paterson HE, De Meillon B. Isolation of Spondweni virus  
 177 from four species of culicine mosquitoes and a report of two laboratory infections with the virus.  
 178 South African Medical Journal. 1961;35:647-50. Epub 1961/08/05.
- 179 14. Henderson BE, Cheshire PP, Kirya GB, Lule M. Immunologic studies with yellow fever  
 180 and selected African group B arboviruses in rhesus and vervet monkeys. The American Journal  
 181 of Tropical Medicine and Hygiene. 1970;19(1):110-8.
- 182 15. Foy BD, Kobylinski KC, Chilson Foy JL, Blitvich BJ, Travassos da Rosa A, Haddow  
 183 AD, et al. Probable non-vector-borne transmission of Zika virus, Colorado, USA. Emerging  
 184 Infectious Diseases. 2011;17(5):880-2.
- 185 16. Wolfe MS, Calisher CH, McGuire K. Spondweni virus infection in a foreign resident of  
 186 Upper Volta. Lancet. 1982;2(8311):1306-8. Epub 1982/12/11.
- 187 17. Heang V, Yasuda CY, Sovann L, Haddow AD, Travassos da Rosa AP, Tesh RB, et al.  
 188 Zika virus infection, Cambodia, 2010. Emerging Infectious Diseases. 2012;18(2):349-51.

- 189 18. Oehler E, Watrin L, Larre P, Leparc-Goffart I, Lastere S, Valour F, et al. Zika virus  
190 infection complicated by Guillain-Barre syndrome--case report, French Polynesia, December  
191 2013. Euro surveillance. 2014;19(9).
- 192 19. Besnard M, Lastere S, Teissier A, Cao-Lormeau V, Musso D. Evidence of perinatal  
193 transmission of Zika virus, French Polynesia, December 2013 and February 2014. Euro  
194 surveillance. 2014;19(13).
- 195